arXiv:nucl-th/0403074v2 2 Apr 2004

Indication of quark deconfinement and evidence for a Hubble flow in 130 and 200 GeV Au+Au collisions

M. Csanád[1], **T. Csörgő**[2], **B. Lörstad**[3] **and A. Ster**[2]
[1]Dept. Atomic Phys., ELTE, H-1117 Budapest, Pázmány P. 1/a, Hungary
[2]MTA KFKI RMKI, H - 1525 Budapest 114, P.O.Box 49, Hungary
[3]Dept. Physics, University of Lund, S - 22362 Lund, Sweden

Abstract. Buda-Lund hydro model fits are compared to BRAHMS, PHENIX, PHOBOS and STAR data on identified particle spectra, two-particle Bose-Einstein or HBT correlations, charged particle pseudorapidity distributions and pseudorapidity as well as p_t dependent elliptic flow in $\sqrt{s_{\rm NN}} = 130$ and 200 GeV Au+Au collisions at RHIC. Preliminary results indicate that 7/8-th of the particle emitting volume is rather cold, with surface temperature of 105 MeV, but the temperature has a distribution and the most central 1/8-th of the volume is superheated to $T(x) > T_c = 172 \pm 3$ MeV.

1. Introduction

The Buda-Lund hydro model [1] is successful in describing the BRAHMS, PHENIX, PHOBOS and STAR data on identified single particle spectra and the transverse mass dependent Bose-Einstein or HBT radii as well as the pseudorapidity distribution of charged particles in Au + Au collisions both at $\sqrt{s_{\rm NN}} = 130$ GeV [2] and at $\sqrt{s_{\rm NN}} = 200$ GeV [3]. Recently, Fodor and Katz calculated the phase diagram of lattice QCD at finite net baryon density [4]. Their results, obtained with light quark masses four times heavier than the physical value, indicated that in the $0 \leq \mu_B \leq 300$ MeV region the transition from confined to deconfined matter is not a first or second order phase-transition, but a cross-over with a nearly constant critical temperature, $T_c = 172 \pm 3$ MeV. The result of the Buda-Lund fits to Au+Au data, both at $\sqrt{s_{\rm NN}} = 130$ and 200 GeV, indicate the existence of a very hot region. The temperature distribution $T(x)$ of this region is characterized with a central temperature T_0, found to be greater than the critical value calculated from lattice QCD: $T_0 > T_c$ [5]. The Buda-Lund fits thus indicate quark deconfinement in Au + Au collisions at RHIC. The observation of a superheated center in Au+Au collisions at RHIC is confirmed by the analysis of p_t and η dependence of the elliptic flow [5], measured by the PHENIX [8] and PHOBOS collaborations [9, 10]. A similar analysis of Pb+Pb collisions at CERN SPS energies yields central temperatures lower than the critical value, $T_0 < T_c$ [6, 7].

Here we summarize the Buda-Lund fit results as detailed in refs. [2, 3, 5, 6]. See these papers for definitions and discussion of the results as well as more detailed references.

2. Buda-Lund fit results to central Au+Au data at $\sqrt{s_{\rm NN}} = 130$ and 200 GeV



Figure 1. The upper four panels show a simultaneous Buda-Lund fit to 0-5(6) % central Au+Au data at $\sqrt{s_{NN}} = 130$ GeV, refs. [11, 12, 13, 14, 15]. The lower four panels show similar fits to 0-30 % central Au+Au data at $\sqrt{s_{\rm NN}} = 200$ GeV, refs. [16, 17, 18]. The fit parameters are summarized in Table 1.

Analyzing the fit parameters (Table 1) we find that $T_0 > T_c$ by more than 5σ in case of the $0-5(6)\%$ most central Au+Au data at $\sqrt{s_{NN}} = 130$ GeV. We interpret this as an indication of quark deconfinement. In case of the less central $(0-30\%)$ Au+Au data at $\sqrt{s_{NN}} = 200$ GeV, with an improved analysis we find $T_0 > T_c$ by 2σ, not a significant difference. We interpret this as a possible hint for quark deconfinement. In both cases, the flow profile within errors coincides with the three-dimensional Hubble flow, $u^\mu = x^\mu/\tau$. For details, see refs. [7, 19]. For similar results, see refs. [20, 21, 22].

3. Buda-Lund results for the elliptic flow in Au+Au data at RHIC



Figure 2. In the top panels, the ellipsoidally symmetric Buda-Lund hydro calculations are compared to PHENIX identified $v_2(p_t)$ data at 200 GeV [8] and PHOBOS $v_2(\eta)$ data at 130 and 200 GeV [9, 10]. On the top right panel, we added a constant non-flow parameter of 0.02 to the calculated values of v_2. In the lower panels, the flow coefficients, v_n-s are shown for $n = 1$ ... 6 as a function of transverse momentum at midrapidity, as well as a function of pseudorapidity, with integrated p_t.

The model is evaluated in all panels with the following parameters: $T_0 = 210$ MeV, $T_s = 105$ MeV, $X_f = 8.6$ fm, $Y_f = 10.5$ fm, $Z_f = 17.5$ fm, $\dot{X}_f = 0.57$, $\dot{Y}_f = 0.45$, $\dot{Z}_f = 2.4$, $\tau_0 = 7$ fm/c, $\vartheta = 0.09$, $\mu_{0,\pi} = 70$ MeV, $\mu_{0,K} = 210$ MeV and $\mu_{0,p} = 315$ MeV. T_0 is the temperature of the center, while T_s is that of the surface, (X_f, Y_f, Z_f) are the principal axes and $(\dot{X}_f, \dot{Y}_f, \dot{Z}_f)$ the principal expansion rates of the ellipsoid at the freeze-out proper-time τ_0. For more details, see pages 3-6 of ref. [5].

Table 1, Figures 1 and 2 indicate that the Buda-Lund hydro model works well at both RHIC energies: it gives a good quality description of the transverse mass dependence of the HBT radii as well as identified particle spectra and elliptic flow, see refs. [1, 23, 24, 25] for further details.

The temperature was estimated to be above T_c in the most central 1/8th of the expanding ellipsoid [5], similarly to a fireball that is heated from inside. We interpret this result as a confirmation of the quark deconfinement and the cross-over like transition found in Buda-Lund fits to central Au+Au collisions at $\sqrt{s_{\rm NN}} = 130$ and 200 GeV.

This work was supported by following grants: OTKA T034269, T038406, the OTKA-MTA-NSF grant INT0089462, the NATO PST.CLG.980086 grant and the exchange program of the Hungarian and Polish Academy of Sciences.

Buda-Lund parameter	Au+Au 200 GeV	Au+Au 130 GeV
T_0 [MeV]	196 ± 13	214 ± 7
T_e [MeV]	117 ± 12	102 ± 11
μ_B [MeV]	61 ± 52	77 ± 38
R_G [fm]	13.5 ± 1.7	28.0 ± 5.5
R_s [fm]	12.4 ± 1.6	8.6 ± 0.4
$\langle u_t' \rangle$	1.6 ± 0.2	1.0 ± 0.1
τ_0 [fm/c]	5.8 ± 0.3	6.0 ± 0.2
$\Delta\tau$ [fm/c]	0.9 ± 1.2	0.3 ± 1.2
$\Delta\eta$	3.1 ± 0.1	2.4 ± 0.1
χ^2/NDF	114 / 208	158.2 / 180

Table 1. Buda-Lund hydro model v1.5 source parameters, corresponding to Figs. 1 and 2. The errors on the parameters are preliminary, as point-to-point and normalization errors are added in quadrature when evaluating χ^2.

[1] T. Csörgő and B. Lörstad, Phys. Rev. C **54** (1996) 1390
[2] M. Csanád, T. Csörgő, B. Lörstad, A. Ster, Act. Phys. Pol. B**35**, 191 (2004), nucl-th/0311102.
[3] M. Csanád, T. Csörgő, B. Lörstad and A. Ster, nucl-th/0402037.
[4] Z. Fodor and S. D. Katz, JHEP **0203** (2002) 014
[5] M. Csanád, T. Csörgő and B. Lörstad, nucl-th/0310040.
[6] A. Ster, T. Csörgő and B. Lörstad, Nucl. Phys. A **661**, 419 (1999) hep-ph/9907338.
[7] T. Csörgő, Heavy Ion Phys. **15**, 1 (2002)
[8] S. S. Adler *et al.* [PHENIX Collaboration], nucl-ex/0305013.
[9] B. B. Back *et al.* [PHOBOS Coll.], Phys. Rev. Lett. **89**, 222301 (2002)
[10] S. Manly *et al.* [PHOBOS Collaboration], Nucl. Phys. A**715**, 611 (2003)
[11] I. G. Bearden *et al.* [BRAHMS Collaborations], Phys. Lett. B **523** (2001) 227
[12] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88** (2002) 242301
[13] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88** (2002) 192302
[14] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **87** (2001) 102303
[15] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **87** (2001) 082301
[16] I. G. Bearden *et al.* [BRAHMS Collaboration], Phys. Rev. Lett. **88** (2002) 202301
[17] S. S. Adler *et al.* [PHENIX Collaboration], nucl-ex/0307022.
[18] S. S. Adler *et al.* [PHENIX Collaboration], nucl-ex/0401003.
[19] http://www.kfki.hu/˜csorgo/budalund/
[20] F. Retiere and M. A. Lisa, arXiv:nucl-th/0312024.
[21] W. Broniowski and W. Florkowski, Phys. Rev. Lett. **87** (2001) 272302 [arXiv:nucl-th/0106050].
[22] W. Broniowski, A. Baran and W. Florkowski, AIP Conf. Proc. **660** (2003) 185
[23] T. Csörgő and A. Ster, Heavy Ion Phys. **17** (2003) 295
[24] T. Csörgő and J. Zimányi, Heavy Ion Phys. **17**, 281 (2003)
[25] T. Csörgő, L. Csernai, Y. Hama, T. Kodama, nucl-th/0306004.